

PMRA Approved Label
2020-1214
2021-06-21

GROUP	3	11	FUNGICIDE
-------	---	----	-----------

VELTYMA®

FUNGICIDE

For use in canola, corn (field, pop, sweet, seed), dried shelled pea and bean (Crop Subgroup 6C), flax, mustard, potato, rapeseed, soybean, sugar beet, and wheat (all types)

Suspension Concentrate

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS: Mefentrifluconazole 200 g/L
Pyraclostrobin 200 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00111%, 2-methyl-4-isothiazolin-3-one at 0.00037%, 1,2-benzisothiazolin-3-one at 0.0400% and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

Warning: contains the allergen soy

REGISTRATION NO. 34166 PEST CONTROL PRODUCTS ACT



CAUTION POISON

SKIN IRRITANT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 0.1 L – Bulk

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

VELTYMA® is a registered trademark of BASF Agro B.V. Arnhem (NL), Zweigniederlassung Freienbach, used with permission by BASF Canada Inc.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. May irritate the skin. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours .
6. Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Coveralls and gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. **DO NOT** contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
10. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, non-target terrestrial plants, and small wild mammals. Observe spray buffer zones specified under DIRECTIONS FOR USE

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Mefentrifluconazole is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing mefentrifluconazole during the previous season.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For Non-Returnable or Non-refillable Containers

Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

GROUP	3	11	FUNGICIDE
-------	---	----	-----------

VELTYMA®

FUNGICIDE

For use in canola, corn (field, pop, sweet, seed), dried shelled pea and bean (Crop Subgroup 6C), flax, mustard, potato, rapeseed, soybean, sugar beet, and wheat (all types)

Suspension Concentrate

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS: Mefentrifluconazole 200 g/L
Pyraclostrobin 200 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00111%, 2-methyl-4-isothiazolin-3-one at 0.00037%, 1,2-benzisothiazolin-3-one at 0.0400% and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

Warning: contains the allergen soy

REGISTRATION NO. 34166 PEST CONTROL PRODUCTS ACT



CAUTION POISON

SKIN IRRITANT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 0.1 L – Bulk

BASF Canada Inc.
100 Milverton Drive, 5th Floor
Mississauga, Ontario
L5R 4H1
1-877-371-2273

VELTYMA® is a registered trademark of BASF Agro B.V. Arnhem (NL), Zweigniederlassung Freienbach, used with permission by BASF Canada Inc.

GENERAL INFORMATION

This package contains **VELTYMA**, a broad-spectrum fungicide. The active ingredients in **VELTYMA** belong to two classes of fungicides; mefentrifluconazole belongs to the triazole (DMI) group and pyraclostrobin belongs to the strobilurin (QoI) group of fungicides.

VELTYMA inhibits spore germination, mycelial growth and sporulation of the fungus on the leaf surface. **VELTYMA** can be applied in either pre- or post-infection situations. However, optimum disease control is achieved when **VELTYMA** is applied preventatively in a regularly scheduled spray program and is used in a rotation program with other fungicides.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. May irritate the skin. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours .
6. Wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Coveralls and gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. **DO NOT** contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
10. Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, non-target terrestrial plants, and small wild mammals. Observe spray buffer zones specified under DIRECTIONS FOR USE

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Mefentrifluconazole is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing mefentrifluconazole during the previous season.

DIRECTIONS FOR USE (See specific sections for each crop group)

<p>All Crops</p> <p>Apply recommended rates of VELTYMA as instructed in the following series of crop application rate tables. Apply VELTYMA with spray equipment as specified. Equipment should be checked frequently for proper calibration.</p> <p>Apply before disease development. See crop-specific requirements for exceptions.</p> <p>Select the use rate based on relative disease pressure. Use the high rate if weather conditions are favorable for disease development (i.e. high humidity/moisture) and/or when risk for disease development is high (i.e. narrow host rotation with disease history and high potential for inoculum).</p> <p>DO NOT make more than two (2) sequential applications of VELTYMA targeting the same disease before alternating to a labeled fungicide containing a different mode of action.</p>
<p>TANK MIXING</p> <p>In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact BASF Canada Inc. at 1-877-371-2273 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.</p> <p>Follow the most restrictive label directions for each of the tank mix partners including the rates, maximum number of applications, application timing, protective equipment, restrictions and precautions.</p>

Canola, Rapeseed (Brassica spp., Brassica napus), Mustard (Brassica spp., Brassica juncea) including oilseed/condiment mustard Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Blackleg (<i>Leptosphaeria maculans</i>)	0.375 - 0.5	21	1.0
<p>Apply VELTYMA at the 2 to 6-leaf stage to control blackleg. Use the high rate under high disease pressure. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.</p> <p>Canola/rapeseed forage cannot be grazed or fed to livestock.</p>			
Detailed Crop Listing: canola, rapeseed, mustard			

Corn Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Common rust (<i>Puccinia sorghi</i>) Eyespot (<i>Kabatiella zea</i> / <i>Aureobasidium zea</i>) Gray leaf spot (<i>Cercospora zea-maydis</i>) Northern leaf blight (<i>Exserohilum turcicum</i> / <i>Setosphaeria turcica</i>) Tar spot (<i>Phyllachora maydis</i>)	0.5	21	1.0
<p>For optimal disease control, begin applications of VELTYMA prior to disease development. Apply a second time 10-14 days later if disease persists or weather conditions are favourable for disease development.</p>			
Detailed Crop Listing: corn (field, pop, sweet, and seed)			

**Dried Shelled Peas and Beans (Crop Subgroup 6C)
Application Rate Table
(Ground and Aerial Application)**

Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Anthracnose (<i>Colletotrichum lentis</i> ; <i>C. lindemuthianum</i>) Ascochyta blight (<i>Ascochyta pinodes</i> ; <i>A. lentis</i> ; <i>A. fabae</i> , <i>A. rabiei</i>) Mycosphaerella blight (<i>Mycosphaerella pinodes</i>) Powdery mildew (<i>Erysiphe polygoni</i>) Rust (<i>Uromyces appendiculatus</i>)	0.5	30	1.0
Disease Suppressed Downy mildew (<i>Peronospora viciae</i> f. sp. <i>pisii</i>) (dry pea only)			

Apply **VELTYMA** at the beginning of flowering or at the onset of symptoms. Apply a second time 10-14 days later if disease persists, or weather conditions are favourable for disease development.

Detailed Crop Listing: Crop Subgroup 6C: Dried shelled peas and beans (except soybean) including:

Bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin and white sweet lupin);

Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean);

Bean (*Vigna* spp.) (includes adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean);

Broad bean (fava bean) (*Vicia faba*); Chickpea (garbanzo bean) (*Cicer arietinum*); Guar (*Cyamopsis tetragonoloba*); Lablab bean (hyacinth bean) (*Lablab purpureus*); Lentil (*Lens esculenta*); Pea (*Pisum* spp.) (includes field pea); Pigeon pea (*Cajanus cajan*).

Flax Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Pasmo (<i>Septoria linicola</i>)	0.375 - 0.5	21	1.0
Apply VELTYMA beginning at 20-50% flowering to control pasmo. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			

Potato Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Early Blight (<i>Alternaria solani</i>)*	0.5	7	1.0
Black Dot (<i>Colletotrichum coccodes</i>)			
Disease Suppressed Brown spot (<i>Alternaria alternata</i>)			
Begin applications of VELTYMA prior to row closure or when conditions are favorable for disease development (whichever comes first). Apply a second time 7 - 14 days later if disease persists or weather conditions are favourable for disease development. *Includes control of Group 11 resistant biotypes			

Soybean Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval	Maximum per year (L/ha)

		(All Commodities) (days)	
Cercospora Blight & Purple Seed Stain (<i>Cercospora kikuchii</i>)	0.375 – 0.5	21	1.0
Septoria Brown Spot (<i>Septoria glycines</i>)			
Frogeye Leaf Spot (<i>Cercospora sojina</i>)*	0.5		
Pod and stem blight (<i>Diaporthe phaseolorum</i> var. <i>sojai</i> / <i>Phomopsis longicolla</i>)			
<p>Apply VELTYMA prior to disease development when conditions are favorable for disease development. Apply a second time 10-14 days later if disease persist or weather conditions are favourable for disease development.</p> <p>*Includes control of Group 11 resistant biotypes</p>			

**Sugar Beet Application Rate Table
(Ground and Aerial Application)**

Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)
Cercospora leaf spot (<i>Cercospora beticola</i>)*	0.5 – 0.75	21	1.5
<p>Begin applications of VELTYMA prior to disease development. Apply additional applications 14 days later if disease persists or weather conditions are favourable for disease development.</p> <p>*Includes control of Group 11 resistant biotypes</p>			

**Wheat (All Types) Application Rate Table
(Ground and Aerial Application)**

Disease Controlled	Application Rate (L/ha)	PHI; Livestock Grazing and/or Feeding Interval (All Commodities) (days)	Maximum per year (L/ha)

Leaf rust (<i>Puccinia recondita</i>)	0.375 – 0.5	21	1.0
Stripe rust (<i>Puccinia striiformis</i>)			
Septoria leaf blotch (<i>Septoria tritici</i> or <i>Septoria nodorum</i>)			
Tan spot (<i>Pyrenophora tritici-repentis</i>)			
<p>Apply VELTYMA prior to disease development, when conditions are favorable for disease development, or at the onset of disease symptoms. Apply a second time 10-14 days later if disease persists or weather conditions are favourable for disease development.</p> <p>To control foliar leaf diseases, VELTYMA can be applied between stem elongation stage (GS 31) to early head emergence (GS 55). To maximize yield in wheat, it is important to protect the flag leaf from disease. Therefore the optimum time to apply a single application of VELTYMA is immediately after flag leaf emergence (GS 37-39).</p>			
<p>Detailed Crop Listing: wheat (spring, winter, and durum)</p>			

APPLICATION INSTRUCTIONS

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Ground Application

Apply **VELTYMA** at rates listed in the Application Rate and Timing Tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 100 L/ha and ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

Aerial Application

Apply **VELTYMA** at rates listed in application rate and timing tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 50 L/ha. Ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

Instructions for Aerial Applications

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-877-371-BASF (2273) or obtain technical advice from the distributor or your provincial agricultural representative.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for uses with hand-held application equipment on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:			
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Canola, Rapeseed (<i>Brassica</i> spp., <i>Brassica napus</i>), Mustard (<i>Brassica</i> spp., <i>Brassica juncea</i>) including oilseed/condiment mustard, Corn, Dried shelled peas and beans (Crop Subgroup 6C), Flax, Soybean, Wheat, Potato, Sugarbeets		5	1	1	1
Aerial	Rapeseed (<i>Brassica</i> spp., <i>Brassica napus</i>), Mustard (<i>Brassica</i> spp., <i>Brassica juncea</i>) including oilseed/condiment mustard, Corn, Dried shelled peas and beans (Crop Subgroup 6C), Flax, Soybean, Wheat	Fixed wing	125	10	5	0

	Potato	Rotary wing	100	10	2	0
		Fixed wing	150	10	5	0
	Sugarbeets	Rotary wing	125	10	2	0
		Fixed wing	125	10	5	0
		Rotary wing	95	10	2	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **VELTYMA** contains Group 3 and 11 fungicides. Any fungal population may contain individuals naturally resistant to **VELTYMA** and other Group 3 or 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same locations. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of **VELTYMA** with a fungicide from a different mode of action other than Group 3 or 11 that controls the same pathogens. **DO NOT** exceed the total number of applications of **VELTYMA** per season as stated in the Crop-Specific Restrictions and Limitations.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Use tank mixtures with fungicide/bactericides from a different group that is effective on the target pathogen when such use is permitted.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify BASF Canada if reduced sensitivity of the pathogen to **VELTYMA** is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate.

Discontinue use of this product, and switch to another fungicide with a different site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information, contact BASF Canada Inc. at 1-877-371-2273 or www.agsolutions.ca.

MIXING

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of **VELTYMA** to the tank.
4. Add the required amount of the tank mix partner.
5. Add the recommended amount of adjuvant.
6. Continue agitation while filling the remainder of the spray tank .
7. After use, clean the spray tank according to PRECAUTIONS.

RESTRICTIONS AND LIMITATIONS

1. All crops on this label can be grazed or fed to livestock, except for canola (rapeseed) forage. For livestock feeding and grazing, observe the minimum preharvest intervals specified for each crop under the Crop-Specific Table.
2. Crop Rotation Restrictions: All crops on this label may be planted immediately after the last application. Crops not on this label must not be planted within 1 month after the last application.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For Non-Returnable or Non-Refillable Containers

Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.